

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use XCOPRI safely and effectively. See full prescribing information for XCOPRI.

XCOPRI® (cenobamate tablets), for oral use, CV
Initial U.S. Approval: 2020

RECENT MAJOR CHANGES

Dosage and Administration, Administration Instructions (2.3) 4/2024

INDICATIONS AND USAGE

XCOPRI is indicated for the treatment of partial-onset seizures in adult patients. (1)

DOSAGE AND ADMINISTRATION

- The recommended initial dosage of XCOPRI is 12.5 mg once daily, titrated to the recommended maintenance dosage of 200 mg once daily. The recommended titration schedule should not be exceeded. The maximum dosage is 400 mg once daily. (2.1)
- Hepatic impairment: For patients with mild or moderate hepatic impairment, the maximum recommended dosage is 200 mg once daily. (2.2, 8.7, 12.3)
- XCOPRI can be taken whole or the tablets can be crushed. The crushed tablet can be mixed with water and either administered by mouth as an oral suspension or administered via a nasogastric tube. (2.3)

DOSAGE FORMS AND STRENGTHS

- Tablets: 12.5 mg, 25 mg, 50 mg, 100 mg, 150 mg, and 200 mg. (3)

CONTRAINDICATIONS

- Hypersensitivity to cenobamate or any of the inactive ingredients in XCOPRI. (4)
- Familial Short QT syndrome. (4)

WARNINGS AND PRECAUTIONS

- Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multi-Organ Hypersensitivity*: Discontinue if no alternate etiology. (5.1)
- QT Shortening*: Use caution when administering XCOPRI with other drugs that shorten the QT interval (5.2)
- Suicidal Behavior and Ideation*: Monitor patients for suicidal behavior and ideation. (5.3)
- Neurological Adverse Reactions*: Monitor for somnolence and fatigue and advise patients not to drive or operate machinery until they have gained

sufficient experience on XCOPRI. Concomitant use with other CNS depressants or alcohol may have additive effects. (5.4)

- Withdrawal of Antiepileptic Drugs*: XCOPRI should be gradually withdrawn to minimize the potential of increased seizure frequency. (5.5)

ADVERSE REACTIONS

The most common adverse reactions in patients receiving XCOPRI (at least 10% for XCOPRI and more frequently than placebo) include somnolence, dizziness, fatigue, diplopia, and headache. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact SK Life Science, Inc. at 1-866-657-5574 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Phenytoin: Gradually decrease phenytoin dosage by up to 50% (7.1)
- Phenobarbital and Clobazam: Reduce dosage as needed when used concomitantly with XCOPRI. (7.1)
- Lamotrigine, Carbamazepine: Increase dosage as needed when used concomitantly with XCOPRI. (7.1)
- CYP2B6 and CYP3A Substrates: Increase dosage as needed when used concomitantly with XCOPRI. (7.1)
- CYP2C19 Substrates: Reduce dosage as needed when used concomitantly with XCOPRI. (7.1)
- Oral Contraceptives: Effectiveness of hormonal oral contraceptives may be reduced when administered concomitantly with XCOPRI. Women should use additional or alternative non-hormonal birth control. (7.1)

USE IN SPECIFIC POPULATIONS

- Pregnancy: Based on animal data, may cause fetal harm. (8.1)
- Renal Impairment: Use with caution and dosage reduction may be considered in patients with mild to moderate (CL_{cr} 30 to < 90 mL/min) and severe (CL_{cr} < 30 mL/min) renal impairment. Use not recommended in end-stage renal disease (CL_{cr} < 15 mL/min) undergoing dialysis. (8.6)
- Hepatic Impairment: Use with caution in patients with mild to moderate hepatic impairment; lower maximum dosage and additional dosage reduction may be considered. Use of XCOPRI in patients with severe hepatic impairment is not recommended. (2.2, 8.7, 12.3)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 4/2024

FULL PRESCRIBING INFORMATION: CONTENTS*

1	INDICATIONS AND USAGE	8.3	Females and Males of Reproductive Potential
2	DOSAGE AND ADMINISTRATION	8.4	Pediatric Use
2.1	General Dosing Recommendations	8.5	Geriatric Use
2.2	Dosage Modifications in Patients with Hepatic Impairment	8.6	Patients with Renal Impairment
2.3	Administration Instructions	8.7	Patients with Hepatic Impairment
2.4	Discontinuation of XCOPRI	9	DRUG ABUSE AND DEPENDENCE
3	DOSAGE FORMS AND STRENGTHS	9.1	Controlled Substance
4	CONTRAINDICATIONS	9.2	Abuse
5	WARNINGS AND PRECAUTIONS	9.3	Dependence
5.1	Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multiorgan Hypersensitivity	10	OVERDOSAGE
5.2	QT Shortening	11	DESCRIPTION
5.3	Suicidal Behavior and Ideation	12	CLINICAL PHARMACOLOGY
5.4	Neurological Adverse Reactions	12.1	Mechanism of Action
5.5	Withdrawal of Antiepileptic Drugs	12.2	Pharmacodynamics
6	ADVERSE REACTIONS	12.3	Pharmacokinetics
6.1	Clinical Trials Experience	13	NONCLINICAL TOXICOLOGY
6.2	Postmarketing Experience	13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
7	DRUG INTERACTIONS	14	CLINICAL STUDIES
7.1	Effect of XCOPRI on Other Drugs	16	HOW SUPPLIED/STORAGE AND HANDLING
7.2	Drugs that Shorten the QT Interval	16.1	How Supplied
7.3	CNS Depressants and Alcohol	16.2	Storage and Handling
8	USE IN SPECIFIC POPULATIONS	17	PATIENT COUNSELING INFORMATION
8.1	Pregnancy	*Sections or subsections omitted from the full prescribing information are not listed.	
8.2	Lactation		

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

XCOPRI is indicated for the treatment of partial-onset seizures in adult patients.

2 DOSAGE AND ADMINISTRATION

2.1 General Dosing Recommendations

Monotherapy and Adjunctive Therapy

XCOPRI is administered orally once daily with or without food. The recommended dosage and titration, which should not be exceeded because of the potential for serious adverse reactions [*see Warnings and Precautions (5.2)*], is included in Table 1.

Table 1: Recommended Dosage for Partial-Onset Seizures in Adults

Initial Dosage	
Week 1 and 2	12.5 mg once daily
Titration Regimen	
Week 3 and 4	25 mg once daily
Week 5 and 6	50 mg once daily
Week 7 and 8	100 mg once daily
Week 9 and 10	150 mg once daily
Maintenance Dosage	
Week 11 and thereafter	200 mg once daily
Maximum Dosage	
If needed based on clinical response and tolerability, dose may be increased above 200 mg by increments of 50 mg once daily every two weeks to 400 mg.	400 mg once daily

2.2 Dosage Modifications in Patients with Hepatic Impairment

For patients with mild to moderate (Child-Pugh Class A to B) hepatic impairment, the maximum recommended dosage is 200 mg once daily [*see Use in Specific Populations (8.7)*]. XCOPRI is not recommended for use in patients with severe (Child-Pugh Class C) hepatic impairment [*see Clinical Pharmacology (12.3)*].

2.3 Administration Instructions

XCOPRI can be taken whole or the tablets can be crushed. The crushed tablet can be mixed with water and either administered by mouth as an oral suspension or administered via a nasogastric tube, as described below [*see Clinical Pharmacology (12.3)*].

Administration of Crushed Tablets by Mouth as Oral Suspension

1. Crush the appropriate number of tablet(s) for the prescribed dose.
2. In a cup, combine the crushed tablet(s) and 25 mL of water.
3. Swirl to suspend the crushed tablet(s).
4. Drink the suspension immediately. Do not store the tablet-water mixture for later use.
5. To ensure no tablet residue is left in the container, rinse the container with 25 mL of water and drink.
6. Visually confirm that no particles are left in the container. If particles remain, repeat step 5.

Administration of Crushed Tablets via Nasogastric (NG) Tube

1. Crush the appropriate number of tablet(s) for the prescribed dose.
2. In an appropriate container, combine the crushed tablet(s) and 25 mL of water.
3. Swirl to suspend the crushed tablet(s).
4. Ensuring no particles are left in the container, instill the suspension with a syringe into the NG tube.
5. Refill the catheter-tip syringe again with 10 mL of water, swirl gently, and administer.
6. Visually confirm that no particles are left in the syringe. If particles remain, repeat step 5.

2.4 Discontinuation of XCOPRI

If XCOPRI is discontinued, the dosage should be gradually reduced over a period of at least 2 weeks, unless safety concerns require abrupt withdrawal [see *Warnings and Precautions (5.1, 5.5)*].

3 DOSAGE FORMS AND STRENGTHS

XCOPRI tablets are available in the following strengths, shapes, colors, and tablet markings (Table 2).

Table 2: XCOPRI Tablet Presentations

Tablet Strength	Tablet Color/Shape	Tablet Markings
12.5 mg	Uncoated round white to off-white tablets	SK on one side and 12 on the other side
25 mg	Film coated round brown tablets	SK on one side and 25 on the other side
50 mg	Film coated round yellow tablets	SK on one side and 50 on the other side
100 mg	Film coated round brown tablets	SK on one side and 100 on the other side
150 mg	Film coated round light orange tablets	SK on one side and 150 on the other side
200 mg	Film coated modified oval light orange tablets	SK on one side and 200 on the other side

4 CONTRAINDICATIONS

XCOPRI is contraindicated in patients with:

- Hypersensitivity to cenobamate or any of the inactive ingredients in XCOPRI [*see Warnings and Precautions (5.1) and Description (11)*]
- Familial Short QT syndrome [*see Warnings and Precautions (5.2)*]

5 WARNINGS AND PRECAUTIONS

5.1 Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multiorgan Hypersensitivity

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS), also known as multiorgan hypersensitivity, has been reported in patients taking XCOPRI. DRESS has occurred, including one fatality, when XCOPRI was titrated rapidly (weekly or faster titration). No cases of DRESS were reported in an open-label safety study of 1339 partial-onset seizure patients when XCOPRI was initiated at 12.5 mg once daily and titrated every two weeks. This finding does not establish that the risk of DRESS is prevented by a slower titration; however, XCOPRI should be initiated at 12.5 mg once daily and titrated every two weeks [*see Dosage and Administration (2.2)*]. DRESS typically, although not exclusively, presents with fever, rash, lymphadenopathy, and/or facial swelling, in association with other organ system involvement, such as hepatitis, nephritis, hematological abnormalities, myocarditis, or myositis sometimes resembling an acute viral infection. Eosinophilia is often present. This disorder is variable in its expression, and other organ systems not noted here may be involved. It is important to note that early manifestations of hypersensitivity, such as fever or lymphadenopathy, may be present even though rash is not evident. If such signs or symptoms are present, the patient should be evaluated immediately. XCOPRI should be discontinued immediately and not restarted if an alternative etiology for the signs or symptoms cannot be established [*see Contraindications (4)*].

5.2 QT Shortening

In a placebo-controlled study of the QT interval, a higher percentage of subjects who took XCOPRI (31% at 200 mg and 66% at 500 mg) had a QT shortening of greater than 20 msec compared to placebo (6-17%). Reductions of the QTc interval below 300 msec were not observed [*see Clinical Pharmacology (12.2)*]. Familial Short QT syndrome is associated with an increased risk of sudden death and ventricular arrhythmias, particularly ventricular fibrillation. Such events in this syndrome are believed to occur primarily when the corrected QT interval falls below 300 msec. Nonclinical data also indicate that QT shortening is associated with ventricular fibrillation. Patients with Familial Short QT syndrome should not be treated with XCOPRI [*see Contraindications (4)*]. Caution should be used when administering XCOPRI and other drugs that shorten the QT interval as there may be a synergistic effect on the QT interval that would increase the QT shortening risk.

5.3 Suicidal Behavior and Ideation

Antiepileptic drugs (AEDs), including XCOPRI, increase the risk of suicidal thoughts or behavior in patients taking these drugs for any indication. Patients treated with any AED for any indication

should be monitored for the emergence or worsening of depression, suicidal thoughts or behavior, and/or any unusual changes in mood or behavior.

Pooled analyses of 199 placebo-controlled clinical trials (mono- and adjunctive therapy) of 11 different AEDs showed that patients randomized to one of the AEDs had approximately twice the risk (adjusted Relative Risk 1.8, 95% CI:1.2, 2.7) of suicidal thinking or behavior compared to patients randomized to placebo. In these trials, which had a median treatment duration of 12 weeks, the estimated incidence rate of suicidal behavior or ideation among 27,863 AED-treated patients was 0.43%, compared to 0.24% among 16,029 placebo-treated patients, representing an increase of approximately one case of suicidal thinking or behavior for every 530 patients treated. There were four suicides in drug-treated patients in the trials and none in placebo-treated patients, but the number is too small to allow any conclusion about drug effect on suicide.

The increased risk of suicidal thoughts or behavior with AEDs was observed as early as one week after starting drug treatment with AEDs and persisted for the duration of treatment assessed. Because most trials included in the analysis did not extend beyond 24 weeks, the risk of suicidal thoughts or behavior beyond 24 weeks could not be assessed.

The risk of suicidal thoughts or behavior was generally consistent among drugs in the data analyzed. The finding of increased risk with AEDs of varying mechanisms of action and across a range of indications suggests that the risk applies to all AEDs used for any indication. The risk did not vary substantially by age (5-100 years) in the clinical trials analyzed.

Table 3 shows absolute and relative risk by indication for all evaluated AEDs.

Table 3: Risk of Suicidal Thoughts or Behaviors by Indication for Antiepileptic Drugs in the Pooled Analysis

Indication	Placebo Patients with Events Per 1000 Patients	Drug Patients with Events Per 1000 Patients	Relative Risk: Incidence of Events in Drug Patients/Incidence in Placebo Patients	Risk Differences: Additional Drug Patients with Events Per 1000 Patients
Epilepsy	1.0	3.4	3.5	2.4
Psychiatric	5.7	8.5	1.5	2.9
Other	1.0	1.8	1.9	0.9
Total	2.4	4.3	1.8	1.9

The relative risk for suicidal thoughts or behavior was higher in clinical trials in patients with epilepsy than in clinical trials in patients with psychiatric or other conditions, but the absolute risk differences were similar for epilepsy and psychiatric indications.

Anyone considering prescribing XCOPRI or any other AED must balance this risk with the risk of untreated illness. Epilepsy and many other illnesses for which AEDs are prescribed are themselves associated with morbidity and mortality and an increased risk of suicidal thoughts and behavior. Should suicidal thoughts and behavior emerge during treatment, the prescriber needs to

consider whether the emergence of these symptoms in any given patient may be related to the illness being treated.

5.4 Neurological Adverse Reactions

Somnolence and Fatigue

XCOPRI causes dose-dependent increases in somnolence and fatigue-related adverse reactions (somnolence, fatigue, asthenia, malaise, hypersomnia, sedation, and lethargy) [see *Adverse Reactions (6.1)*]. In Study 1 and Study 2, 31% of patients randomized to receive XCOPRI at 100 mg/day, 36% of patients randomized to receive XCOPRI at 200 mg/day, and 57% of patients randomized to receive XCOPRI at 400 mg/day reported at least one of these adverse reactions, compared to 19% of patients who received placebo. Somnolence and fatigue-related adverse reactions were serious in 0.4% of XCOPRI-treated patients compared to no patients who received placebo and led to discontinuation in 2% of XCOPRI-treated patients compared to 1% of patients who received placebo.

Dizziness and Disturbance in Gait and Coordination

XCOPRI causes dose-dependent adverse reactions related to dizziness and disturbance in gait and coordination (dizziness, vertigo, balance disorder, ataxia, nystagmus, gait disturbance, and abnormal coordination) [see *Adverse Reactions (6.1)*]. In Study 1 and Study 2, 21% of patients randomized to receive XCOPRI at 100 mg/day, 31% of patients randomized to receive XCOPRI at 200 mg/day, and 52% of patients randomized to receive XCOPRI at 400 mg/day reported at least one of these adverse reactions, compared to 18% of patients who received placebo. Dizziness and disturbance in gait and coordination adverse reactions were serious in 2% of XCOPRI-treated patients compared to no patients who received placebo and led to discontinuation in 5% of XCOPRI-treated patients compared to 1% of patients who received placebo.

Cognitive Dysfunction

XCOPRI causes adverse reactions related to cognitive dysfunction related-events (i.e., memory impairment, disturbance in attention, amnesia, confusional state, aphasia, speech disorder, slowness of thought, disorientation, and psychomotor retardation) [see *Adverse Reactions (6.1)*]. In Study 1 and Study 2, 6% of patients randomized to receive XCOPRI at 100 mg/day, 6% of patients randomized to receive XCOPRI at 200 mg/day, and 9% of patients randomized to receive XCOPRI at 400 mg/day reported at least one of these adverse reactions, compared to 2% of patients who received placebo. No cognitive dysfunction-related events were serious in XCOPRI-treated patients or in patients who received placebo. Cognitive dysfunction related adverse reactions led to discontinuation in 0.4% of XCOPRI-treated patients compared to no patients who received placebo.

Visual Changes

XCOPRI causes adverse reactions related to visual changes including diplopia, blurred vision, and impaired vision [see *Adverse Reactions (6.1)*]. In Study 1 and Study 2, 9% of patients randomized to receive XCOPRI at 100 mg/day, 9% of patients randomized to receive XCOPRI at 200 mg/day, and 18% of patients randomized to receive XCOPRI at 400 mg/day reported at least one of these adverse reactions, compared to 2% of patients who received placebo. No visual change-related events were serious in XCOPRI-treated patients or in patients who received placebo. Visual

change led to discontinuation in 0.5% of XCOPRI-treated patients compared to no patients who received placebo.

Risk Amelioration

Prescribers should advise patients against engaging in hazardous activities requiring mental alertness, such as operating motor vehicles or dangerous machinery, until the effect of XCOPRI is known. Patients should be carefully observed for signs of central nervous system (CNS) depression, such as somnolence and sedation, when XCOPRI is used with other drugs with sedative properties because of potential additive effects.

5.5 Withdrawal of Antiepileptic Drugs

As with most antiepileptic drugs, XCOPRI should generally be withdrawn gradually because of the risk of increased seizure frequency and status epilepticus [see *Dosage and Administration (2.4) and Clinical Studies (14)*]. But if withdrawal is needed because of a serious adverse event, rapid discontinuation can be considered.

6 ADVERSE REACTIONS

The following serious adverse reactions are described in more detail in the Warnings and Precautions section of the labeling:

- Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multiorgan Hypersensitivity [see *Warnings and Precautions (5.1)*]
- QT Shortening [see *Warnings and Precautions (5.2)*]
- Suicidal Behavior and Ideation [see *Warnings and Precautions (5.3)*]
- Neurological Adverse Reactions [see *Warnings and Precautions (5.4)*]
- Withdrawal of Antiepileptic Drugs [see *Warnings and Precautions (5.5)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions and for varying durations, adverse reaction frequencies observed in the clinical trials of a drug cannot be directly compared with frequencies in the clinical trials of another drug and may not reflect the frequencies observed in practice.

In all controlled and uncontrolled trials performed in adult partial-onset seizure patients, XCOPRI was administered as adjunctive therapy to 1944 patients. Of these patients, 1575 were treated for at least 6 months, 710 for at least 12 months, 349 for at least 24 months, and 320 for at least 36 months. A total of 658 patients (442 patients treated with XCOPRI and 216 patients treated with placebo) constituted the safety population in the pooled analysis of placebo-controlled studies in patients with partial-onset seizures (Studies 1 and 2) [see *Clinical Studies (14)*]. The adverse reactions presented in Table 4 are based on this safety population; the median length of treatment in these studies was 18 weeks. Of the patients in those studies, approximately 49% were male, 76% were Caucasian, and the mean age was 39 years.

In Study 1 and Study 2, adverse events occurred in 77% of patients treated with XCOPRI and 68% treated with placebo. Table 4 gives the incidence of adverse reactions that occurred in subjects with partial-onset seizures in any XCOPRI treatment group and for which the incidence was greater than placebo during the controlled clinical trials. The most common adverse reactions that occurred in XCOPRI-treated patients (incidence at least 10% and greater than placebo) were somnolence, dizziness, fatigue, diplopia, and headache.

The discontinuation rates because of adverse events were 11%, 9%, and 21% for patients randomized to receive XCOPRI at doses of 100 mg/day, 200 mg/day, and 400 mg/day, respectively, compared to 4% in patients randomized to receive placebo. The adverse reactions most commonly (1% or greater in any XCOPRI treatment group, and greater than placebo) leading to discontinuation, in descending order of frequency, were ataxia, dizziness, somnolence, diplopia, nystagmus, and vertigo.

Table 4: Adverse Reactions in Pooled Placebo-Controlled Adjunctive Therapy Studies in Patients with Partial-Onset Seizures with XCOPRI Frequency in Any Treatment Arm Greater Than 1% Over Placebo

Adverse Reaction	XCOPRI			Placebo n=216 %
	100mg n = 108 %	200mg n= 223 %	400mg n=111 %	
Cardiac Disorders				
Palpitations	0	0	2	0
Ear and Labyrinth Disorders				
Vertigo	1	1	6	1
Eye Disorders				
Diplopia	6	7	15	2
Vision Blurred	2	2	4	0
Gastrointestinal Disorders				
Nausea	6	6	9	3
Constipation	2	4	8	0
Diarrhea	1	3	5	0
Vomiting	2	4	5	0
Dry Mouth	1	1	3	0
Abdominal Pain	2	2	1	0
Dyspepsia	2	2	0	0
Infections and Infestations				
Nasopharyngitis	2	4	5	3
Pharyngitis	1	2	0	0
Urinary Tract Infection	2	5	0	2
Injury, Poisoning and Procedural Complications				
Head Injury	1	0	2	0
Investigations				
Alanine Aminotransferase Increased*	1	1	4	0

Aspartate Aminotransferase Increased	1	1	3	0
Weight Decreased	2	0	1	0
Metabolism and Nutrition Disorders				
Decreased Appetite	3	1	5	1
Musculoskeletal and Connective Tissue Disorders				
Back Pain	4	2	5	3
Musculoskeletal Chest Pain	2	1	0	0
Nervous System Disorders				
Somnolence	19	22	37	11
Dizziness	18	22	33	15
Fatigue	12	14	24	7
Headache	10	12	10	9
Balance Disorder	3	5	9	1
Gait Disturbance	1	3	8	1
Dysarthria	2	1	7	0
Nystagmus	3	7	6	0
Ataxia	2	3	6	2
Aphasia	2	1	4	0
Asthenia	0	1	3	1
Dysgeusia	2	0	2	0
Memory Impairment	2	1	2	0
Migraine	0	0	2	0
Sedation	1	1	2	0
Tremor	0	3	1	1
Psychiatric Disorders				
Confusional State	2	2	3	0
Euphoric Mood	0	0	2	0
Irritability	1	0	2	0
Suicidal Ideation	2	1	0	0
Renal and Urinary Disorders				
Pollakiuria	0	1	0	0
Reproductive System and Breast Disorders				
Dysmenorrhea	1	2	1	0
Respiratory, Thoracic and Mediastinal Disorders				
Hiccups	0	1	1	0
Dyspnea	0	3	0	0
Skin and Subcutaneous Tissue Disorders				
Pruritus	2	1	0	0
Rash Papular	2	0	0	0

* Reported as an adverse reaction; see Laboratory Abnormalities for ALT changes from collected laboratory values

Laboratory Abnormalities

Hepatic Transaminases

In Study 2, there was a post-baseline elevation of alanine aminotransferase (ALT) to greater than 3 times the upper limit of normal (ULN) in 1 (0.9%) patient treated with 100 mg XCOPRI, 2 (1.8%) patients treated with 200 mg, and 3 (2.7%) patients treated with 400 mg, compared to no patients who took placebo. The maximum ALT elevation was 7.6 times ULN in patients treated with 400 mg XCOPRI.

Potassium

In clinical studies, there was a post-baseline elevation of potassium values greater than 5 meq/L (upper reference range) in patients treated with XCOPRI. In Study 1, there were 17 (17%) patients treated with XCOPRI 200 mg compared to 8 (7%) patients who took placebo with normal baseline potassium values who had at least one post-baseline maximum value greater than 5 meq/L. In Study 2, there was a dose-related distribution where at least one post-baseline potassium value was greater than 5 meq/L, occurring in 8.3%, 9.1%, and 10.8% of the patients treated with XCOPRI 100 mg, 200 mg, and 400 mg, respectively, compared to 5.6% of patients who took placebo. Two patients had a maximum potassium value of 5.9 meq/L.

Other Adverse Reactions

Gastrointestinal disorders: There was an incidence of appendicitis in the overall clinical trial safety population of 2.9 cases of appendicitis/1000 patient-years of exposure that is in excess of the expected background rate in the general population.

Adverse Reactions Based on Gender

No significant gender differences were noted in the incidence of adverse reactions.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post approval use of XCOPRI. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Psychiatric Disorders: Psychosis (hallucinations, delusions/paranoia), hostility, aggression.

Hepatobiliary Disorders: Hepatic failure

7 DRUG INTERACTIONS

7.1 Effect of XCOPRI on Other Drugs

Table 5 summarizes the effect of XCOPRI on other drugs [see *Clinical Pharmacology (12.3)*].

Table 5: Pharmacokinetic Drug Interactions

Drug or Substrate Type	Effect of XCOPRI on Drug or Substrate	Clinical Recommendation
Antiepileptic Drugs		
lamotrigine	↓ plasma concentrations	Because of a potential for reduced efficacy of these drugs, increase the dosage of lamotrigine or carbamazepine, as needed, when used concomitantly with XCOPRI.
carbamazepine	↓ plasma concentrations	
phenytoin	↑ plasma concentrations	Because of a potential 2-fold increase in phenytoin levels, gradually decrease phenytoin dosage by up to 50% as XCOPRI is being titrated.
phenobarbital	↑ plasma concentrations	Because of a potential for an increase in the risk of adverse reactions from these drugs, consider a reduction in dosage of phenobarbital or clobazam, as clinically appropriate, when used concomitantly with XCOPRI.
desmethylclobazam, the active metabolite of clobazam	↑ plasma concentrations	
CYP2B6 Substrates	↓ plasma concentrations	Because of a potential for reduced efficacy of these drugs, increase the dosage of CYP2B6 or CYP3A4 substrates, as needed, when used concomitantly with XCOPRI.
CYP3A Substrates	↓ plasma concentrations	
Oral contraceptives	↓ plasma concentrations	Because of the potential for reduced efficacy of oral contraceptives, women should use additional or alternative non-hormonal birth control while taking XCOPRI.

Drug or Substrate Type	Effect of XCOPRI on Drug or Substrate	Clinical Recommendation
CYP2C19 Substrates	↑ plasma concentrations	Because of a potential for an increase in the risk of adverse reactions from these drugs, consider a reduction in dosage of CYP2C19 substrates, as clinically appropriate, when used concomitantly with XCOPRI.

7.2 Drug that Shorten the QT Interval

XCOPRI can shorten the QT interval; therefore, caution should be used when administering XCOPRI and other drugs that shorten the QT interval [see *Warnings and Precautions (5.2)* and *Clinical Pharmacology (12.2)*].

7.3 CNS Depressants and Alcohol

Concomitant use of XCOPRI with other CNS depressants, including alcohol, may increase the risk of neurological adverse reactions, including sedation and somnolence [see *Warnings and Precautions (5.4)*].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to antiepileptic drugs (AEDs), such as XCOPRI, during pregnancy. Encourage women who are taking XCOPRI during pregnancy to enroll in the North American Antiepileptic Drug (NAAED) Pregnancy Registry by calling the toll-free number 1-888-233-2334 or visiting <http://www.aedpregnancyregistry.org/>.

Risk Summary

There are no adequate data on the developmental risk associated with the use of XCOPRI in pregnant women.

In animal studies, administration of cenobamate during pregnancy or throughout pregnancy and lactation resulted in adverse effects on development (increased embryofetal mortality, decreased fetal and offspring body weights, neurobehavioral and reproductive impairment in offspring) at clinically relevant drug exposures [see *Data*].

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively. The background risk of major birth defects and miscarriage for the indicated population is unknown.

Data

Animal Data

Oral administration of cenobamate (0, 10, 30, or 60 mg/kg/day) to pregnant rats during the period of organogenesis resulted in increased embryofetal mortality, reduced fetal body weights, and incomplete fetal skeletal ossification at the highest dose tested, which was associated with maternal toxicity. There was a small increase in visceral malformations at the high dose; however, teratogenic potential could not be fully evaluated because of the high rate of embryofetal deaths, which resulted in an inadequate number of fetuses examined. Maternal plasma exposure (AUC) at the no-effect dose for adverse effects on embryofetal development (30 mg/kg/day) was less than that in humans at the maximum recommended human dose (MRHD) of 400 mg.

Oral administration of cenobamate (0, 4, 12, or 36 mg/kg/day) to pregnant rabbits during the period of organogenesis resulted in increased embryofetal mortality at the highest dose tested, which was associated with maternal toxicity. Maternal plasma exposure at the no-effect dose (12 mg/kg/day) for adverse effects on embryofetal development was less than that in humans at the MRHD.

When cenobamate (0, 11, 22, or 44 mg/kg/day) was orally administered to female rats throughout pregnancy and lactation, neurobehavioral impairment (learning and memory deficit and increased auditory startle response) was observed in the offspring at all doses and decreased preweaning body weight gain and adverse effects on reproductive function (decreased numbers of corpora lutea, implantations, and live fetuses) were seen in the offspring at the high dose. Maternal plasma exposure at the lowest effect dose (11 mg/kg/day) for adverse effects on pre- and postnatal development was less than that in humans at the MRHD.

8.2 Lactation

Risk Summary

There are no data available on the presence of cenobamate in human milk, the effects on the breastfed infant, or the effects of the drug on milk production. Cenobamate was present in rat milk at concentrations similar to those in maternal plasma.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for XCOPRI and any potential adverse effects on the breastfed infant from XCOPRI or from the underlying maternal condition.

8.3 Females and Males of Reproductive Potential

Contraception

Women of reproductive potential concomitantly using oral contraceptives should use additional or alternative non-hormonal birth control [*see Drug Interactions (7.1) and Clinical Pharmacology (12.3)*].

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

Juvenile Animal Toxicity Data

Cenobamate was administered orally to juvenile rats from postnatal day (PND) 7 to 70. To maintain consistent plasma drug exposures, doses were increased during the dosing period, up to 120 and 80 mg/kg/day in males and females, respectively. Adverse effects included mortality, delayed sexual maturation, neurological (decreased grip strength) and neurobehavioral (learning and memory deficits) impairment, decreased sperm count, decreased brain weight, and ocular histopathology. Recovery from these effects was observed following discontinuation of dosing. Overall, a no-effect dose for adverse effects on postnatal development was not identified. At the lowest doses tested, plasma cenobamate exposures (AUC) were less than that in humans at the maximum recommended human dose (MRHD) of 400 mg.

8.5 Geriatric Use

Clinical studies of XCOPRI did not include sufficient numbers of patients aged 65 and over to determine the safety and efficacy of XCOPRI in the elderly population. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy [*see Clinical Pharmacology (12.3)*].

8.6 Renal Impairment

XCOPRI should be used with caution and dosage reduction may be considered in patients with mild to moderate (CL_{cr} 30 to less than 90 mL/min) and severe (CL_{cr} less than 30 mL/min) renal impairment. Use in patients with end-stage renal disease undergoing dialysis is not recommended [*see Clinical Pharmacology (12.3)*].

8.7 Hepatic Impairment

XCOPRI should be used with caution in patients with mild to moderate (5-9 points on Child-Pugh assessment; Class A or B) hepatic impairment. In these patients, the maximum recommended dosage is 200 mg once daily and additional dosage reduction may be considered [*see Dosage and Administration (2.2) and Clinical Pharmacology (12.3)*]. Use of XCOPRI in patients with severe (10-15 points on Child-Pugh assessment; Class C) hepatic impairment is not recommended.

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

XCOPRI contains cenobamate, a schedule V controlled substance.

9.2 Abuse

Abuse is the intentional, non-therapeutic use of a drug, even once, for its desirable psychological or physiological effects. In a human abuse potential study conducted in recreational sedative

abusers (n=39), single doses of XCOPRI (200 mg and 400 mg) were compared to placebo. XCOPRI at single doses of 400 mg produced responses on positive subjective measures such as “Drug Liking,” “Overall Drug Liking,” “Take Drug Again,” and “Good Drug Effects” that were statistically greater than the responses produced on these measures by placebo. In this study, euphoric mood occurred at greater extent with XCOPRI (400 mg) (8%) than with placebo (0%). Phase 1 multiple ascending dose studies in healthy subjects showed rates of euphoria and feeling drunk of about 3% and disturbance in attention of about 5% in subjects who received supratherapeutic doses of cenobamate, but these adverse events were absent in the placebo group. In Phase 2 and 3 studies in subjects with epilepsy, euphoric mood, confusional state, and sedation occurred at low rates in subjects who received XCOPRI (0.5-2.5%).

9.3 Dependence

Physical dependence is a state that develops as a result of physiological adaptation in response to repeated drug use, manifested by withdrawal signs and symptoms after abrupt discontinuation or a significant dose reduction of a drug. Clinical studies in healthy subjects indicate that XCOPRI may cause physical dependence and lead to a withdrawal syndrome characterized by insomnia, decreased appetite, depressed mood, tremor, and amnesia. XCOPRI should be withdrawn gradually [see *Warnings and Precautions (5.5)*].


10 OVERDOSAGE

There is limited clinical experience with XCOPRI overdose in humans.

There is no specific antidote for overdose with XCOPRI. In the event of overdose, standard medical practice for the management of any overdose should be used. An adequate airway, oxygenation and ventilation should be ensured; monitoring of cardiac rate and rhythm and vital signs is recommended. A certified poison control center should be contacted for updated information on the management of overdose with XCOPRI. There are no data on the removal of XCOPRI using dialysis.

11 DESCRIPTION

The chemical name of XCOPRI (cenobamate) is [(1*R*)-1-(2-Chlorophenyl)-2-(tetrazol-2-yl) ethyl] carbamate. Its molecular formula is C₁₀H₁₀ClN₅O₂ and its molecular weight is 267.67 g/mol. The chemical structure is:



Cenobamate is a white to off-white crystalline powder. It is slightly soluble in aqueous solutions (water 1.7 mg/mL) and has higher solubility in organic solvents like ethanol (209.4 mg/mL).

XCOPRI tablets are for oral administration and contain the following inactive ingredients: colloidal silicon dioxide, lactose monohydrate, magnesium stearate, microcrystalline cellulose, and sodium starch glycolate and film coating agents specified below:

12.5 mg tablets: Not applicable, since 12.5 mg tablets are uncoated.

25 mg and 100 mg tablets: FD&C Blue# 2/indigo carmine aluminum lake, iron oxide red, iron oxide yellow, polyethylene glycol 3350, polyvinyl alcohol-part hydrolyzed, talc, and titanium dioxide.

50 mg tablets: iron oxide yellow, polyethylene glycol 3350, polyvinyl alcohol-part hydrolyzed, talc, and titanium dioxide.

150 mg and 200 mg tablets: iron oxide red, iron oxide yellow, polyethylene glycol 3350, polyvinyl alcohol-part hydrolyzed, talc, and titanium dioxide.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The precise mechanism by which cenobamate exerts its therapeutic effects in patients with partial-onset seizures is unknown. Cenobamate has been demonstrated to reduce repetitive neuronal firing by inhibiting voltage-gated sodium currents. It is also a positive allosteric modulator of the γ -aminobutyric acid (GABA_A) ion channel.

12.2 Pharmacodynamics

Interactions with Alcohol

No clinically significant differences on objective attention, psychomotor performance, and memory tests, in addition to other subjective CNS tests, were observed following concomitant use of XCOPRI and ethanol (preparation of 40% ethanol in orange juice dosed at 0.7 g/kg for males and 0.57 g/kg for females) in healthy subjects.

Cardiac Electrophysiology

In a placebo-controlled QT study in healthy volunteers, dose-dependent shortening of the QTcF interval has been observed with XCOPRI [see *Warnings and Precautions (5.2)*]. The mean $\Delta\Delta$ QTc is -11 [-13, -8] msec for 200 mg once daily and -18 [-22, -15] msec for 500 mg once daily (1.25 times the maximum recommended dosage). A higher percentage of XCOPRI-treated subjects (31% at 200 mg and 66% at 500 mg) had a QT shortening of greater than 20 msec compared to placebo (6-17%). Reductions of the QTc interval below 300 msec were not observed.

12.3 Pharmacokinetics

Cenobamate AUC increases in a greater than dose-proportional manner following single oral doses from 5 to 750 mg (0.0125 to 1.88 times the maximum recommended dosage). Cenobamate C_{max}

increases in a dose proportional manner. Steady-state plasma concentrations are attained after approximately two weeks of once daily dosing.

The pharmacokinetics of cenobamate are similar when used as monotherapy or as adjunctive therapy for the treatment of partial-onset seizures, except plasma cenobamate multiple-dose exposure (C_{max} , AUC) decreased with co-administration of phenytoin by 27-28%.

Absorption

At least 88% of XCOPRI is absorbed following oral administration, with median T_{max} ranging from 1 to 4 hours.

Plasma C_{max} and AUC for XCOPRI crushed tablets mixed in water, administered either orally or through a nasogastric tube, were similar to whole tablets. The median T_{max} for crushed tablets is 0.5 hours.

Effect of Food

No clinically significant differences in cenobamate pharmacokinetics were observed following administration of a high-fat meal (800 -1000 calories with 50% fat).

Distribution

The apparent volume of distribution (V_d/F) of cenobamate after oral administration of XCOPRI is approximately 40-50 L. Plasma protein binding of cenobamate is 60% and independent of concentration *in vitro*. Cenobamate primarily binds with human albumin protein.

Elimination

The apparent terminal half-life of cenobamate is 50-60 hours and apparent oral clearance is approximately 0.45-0.63 L/hour over a dose range from 100 mg/day to 400 mg/day.

Metabolism

Cenobamate is extensively metabolized. The primary metabolic pathways are by glucuronidation via UGT2B7 and to a lesser extent by UGT2B4, and by oxidation via CYP2E1, CYP2A6, CYP2B6, and to a lesser extent by CYP2C19 and CYP3A4/5.

Following administration of radiolabeled cenobamate, unchanged cenobamate accounted for greater than 98% of the total AUC of radioactivity in plasma. Unchanged cenobamate accounted for 6.8% of the dose which was mainly excreted in the urine (6.4%).

Excretion

Following administration of radiolabeled cenobamate, a mean of 93.0% of the total radioactive dose was recovered in urine (87.8%) and feces (5.2%). More than 50% of the radioactivity was excreted within 72 hours of dosing.

Specific Populations

No clinically significant differences in the pharmacokinetics of cenobamate were observed based on age based on data from subjects age 18 years to 77 years, sex, or race/ethnicity based on data from subjects categorized as Asian, Black, Caucasian, Hispanic, or Other.

Patients with Renal Impairment

Cenobamate plasma AUC was 1.4 fold to 1.5 fold higher in subjects with mild (CLcr 60 to less than 90 mL/min) and moderate (CLcr 30 to less than 60 mL/min) following a single oral 200 mg dose of XCOPRI compared to healthy controls. In subjects with severe (CLcr less than 30 mL/min) renal impairment, cenobamate plasma AUC did not change significantly compared to healthy controls following single oral 100 mg dose of XCOPRI [see *Use in Specific Populations (8.6)*]. The effect of hemodialysis on cenobamate pharmacokinetics has not been studied.

Patients with Hepatic Impairment

Cenobamate plasma AUC was 1.9-fold, 2.3-fold, and 4.2-fold higher in subjects with mild (5-6 points, Child-Pugh Class A), moderate (7-9 points, Child-Pugh Class B), and severe (10-15 points, Child-Pugh Class C) hepatic impairment, respectively, following a single oral 200 mg dose of XCOPRI in subjects with mild and moderate hepatic impairment and 100 mg dose of XCOPRI in subjects with severe hepatic impairment compared to matched healthy controls [see *Dosage and Administration (2.2) and Use in Specific Populations (8.7)*].

Drug Interaction Studies

Clinical Studies

Alcohol

No clinically significant pharmacokinetic differences were observed for either cenobamate or alcohol when administered concomitantly.

AEDs

Multiple doses of concomitant XCOPRI 200 mg once daily increased phenytoin mean C_{max} and AUC by 70% and 84%, respectively, and phenobarbital mean C_{max} and AUC by 34% and 37%, respectively [see *Drug Interactions (7.1)*]. Multiple doses of concomitant XCOPRI 200 mg once daily decreased carbamazepine mean C_{max} and AUC each by 23% [see *Drug Interactions (7.1)*].

No clinically significant differences in the pharmacokinetics of the following drugs were observed when used concomitantly with cenobamate: valproic acid, levetiracetam or lacosamide.

Based on population PK analyses, during treatment within the 100-400 mg/day XCOPRI dose range, lamotrigine concentrations are expected to decrease by 21-52% [see *Drug Interactions (7.1)*]; and levetiracetam concentrations are expected to decrease by 4-13%, which is not expected to be clinically significant.

In subjects treated with XCOPRI in Study 1 and Study 2, there was no clear relationship between efficacy and concomitant oxcarbazepine use. As such, the efficacy data from Study 1 and Study 2 do not support the existence of a clinically relevant interaction perpetrated by XCOPRI against oxcarbazepine.

CYP Substrates

Multiple doses of concomitant XCOPRI 200 mg once daily decreased total bupropion (CYP2B6 substrate) mean C_{max} and AUC by 23% and 39%, respectively, and decreased midazolam (CYP3A substrate) mean C_{max} and AUC by 61% and 72%, respectively [see *Drug Interactions (7.1)*]. Multiple doses of concomitant XCOPRI 200 mg once daily increased the omeprazole (CYP2C19 substrate) mean C_{max} and AUC by 83% and 107%, respectively [see *Drug Interactions (7.1)*]. No

clinically significant differences in the pharmacokinetics of warfarin (CYP2C9 substrate) were observed when used concomitantly with cenobamate.

The effects of concomitant AEDs on cenobamate PK

Plasma cenobamate multiple-dose exposure (C_{max} , AUC) decreased with co-administration of phenytoin by 27-28%. However, repeated dosing of valproate, phenobarbital, and carbamazepine did not have any significant impact on plasma cenobamate multiple-dose exposure.

In Vitro Studies

CYP Enzymes

Cenobamate inhibits CYP2B6, CYP2C19, and CYP3A, but cenobamate does not inhibit CYP1A2, CYP2C8, CYP2C9, or CYP2D6.

Cenobamate induces CYP2B6, CYP2C8, and CYP3A4, but cenobamate does not induce CYP1A2, CYP2C9, or CYP2C19.

Transporters Systems

Cenobamate was not a substrate of P-gp, BCRP, OAT1, OAT3, OCT2, MATE1, or MATE2-K, and cenobamate did not inhibit P-gp, OAT1, OCT1, OCT2, OATP1B3, BSEP, OAT3, or OATP1B1.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Oral administration of cenobamate (0, 5, 15, or 35 mg/kg/day) to Tg.rasH2 mice for up to 26 weeks did not result in an increase in tumors. Oral administration of cenobamate (0, 4, 8, or 20 mg/kg/day) to male and female rats for up to 87 or 90 weeks, respectively, did not result in an increase in tumors. Plasma exposure at the highest dose tested in rats was less than that in humans at the maximum recommended human dose (MRHD) of 400 mg/day.

Mutagenesis

Cenobamate was negative for genotoxicity in in vitro (Ames, mouse lymphoma) and in vivo (rat bone marrow micronucleus) assays.

Impairment of Fertility

Oral administration of cenobamate (0, 11, 22, or 44 mg/kg/day) to male and female rats prior to and throughout mating and continuing in females to Gestation Day 6 did not produce adverse effects on fertility, general reproductive performance, or early embryonic development. Plasma exposure (AUC) at the highest dose tested in rats was less than that in humans at the MRHD.

14 CLINICAL STUDIES

The efficacy of XCOPRI for the treatment of partial-onset seizures was established in two multicenter, randomized, double-blind, placebo-controlled studies in adult patients (Study 1 and

Study 2). Patients enrolled in the studies had partial-onset seizures with or without secondary generalization and were not adequately controlled with 1 to 3 concomitant AEDs. During an 8-week baseline period, patients were required to have at least 3 or 4 partial-onset seizures per 28 days on average with no seizure-free period exceeding 3 to 4 weeks. In these studies, patients had a mean duration of epilepsy of approximately 24 years and median baseline seizure frequency of 8.5 seizures per 28 days. More than 80% of patients were taking 2 or more concomitant AEDs.

Study 1 ([NCT01397968](#)) compared doses of XCOPRI 200 mg/day with placebo. Study 2 ([NCT01866111](#)) compared doses of XCOPRI 100 mg/day, 200 mg/day, and 400 mg/day with placebo. Both studies had an 8-week baseline period to establish a baseline seizure frequency, following which patients were randomized to a treatment arm. Patients entered a treatment period consisting of an initial titration phase (6 weeks), and a subsequent maintenance phase (6 weeks for Study 1 and 12 weeks for Study 2). In Study 1, patients were started on a daily dose of 50 mg (a higher starting dose than currently recommended) and subsequently increased by 50 mg/day every two weeks, until the final daily target dose of 200 mg/day was achieved. In Study 2, patients were started on a daily dose of 50 mg (a higher starting dose than currently recommended) and subsequently increased by 50 mg/day every week (a faster titration than currently recommended) until 100 mg/day or 200 mg/day was reached and then increased by 100 mg/day every week in patients randomized to 400 mg/day [see *Dosage and Administration (2.2)*, *Warnings and Precautions (5.1)*, and *Adverse Reactions (6.1)*].

The primary efficacy outcome in Study 1 and Study 2 was the percent change from baseline in seizure frequency per 28 days in the treatment period. Table 6 summarizes the results of the primary endpoint for XCOPRI in each study.

Table 6: Percent Change from Baseline in Seizure Frequency per 28 Days in the Treatment Period (Study 1 and Study 2)

	N	Median Percent Change from Baseline in Seizure Frequency per 28 Days (%) [*]	p-value (Compared to Placebo)
Study 1			
Placebo	108	-21.5	--
XCOPRI 200 mg/day	113	-55.6	< 0.0001 ^{**}
Study 2			
Placebo	106	-24.3	--
XCOPRI 100 mg/day	108	-36.3	0.006 ^{**}
XCOPRI 200 mg/day	109	-55.2	< 0.001 ^{**}
XCOPRI 400 mg/day	111	-55.3	< 0.001 ^{**}

^{*} A negative percent change from baseline in seizure frequency indicates reduction in seizure frequency from baseline.

^{**} Statistically significant compared to placebo

Figure 1 and Figure 2 show the proportion of patients with different percent reductions during the maintenance phase over baseline in Study 1 and Study 2, respectively. Patients in whom the seizure

frequency increased are shown in the left-most column as “worse.” Patients in whom the seizure frequency decreased are shown in the remaining four categories.

Figure 1: Proportion of Patients Exhibiting Different Percent Reductions During the Maintenance Phase over Baseline in Study 1





Figure 2: Proportion of Patients Exhibiting Different Percent Reductions During the Maintenance Phase over Baseline in Study 2



In Study 2, 4 of 102 (4%) patients in the XCOPRI 100 mg/day group, 11 of 98 (11%) patients in the XCOPRI 200 mg/day group, and 20 of 95 (21%) patients in the XCOPRI 400 mg/day group and 1 of 102 (1%) of patients in the placebo group reported no partial seizures during the maintenance phase.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

XCOPRI tablets are supplied in the following configurations:

Bottles; 30 count

Strength	NDC Number	Tablet Description (Color, Shape, Markings)
25 mg	71699-025-30	Film coated round brown tablets with SK on one side and 25 on the other side
50 mg	71699-050-30	Film coated round yellow tablets with SK on one side and 50 on the other side
100 mg	71699-100-30	Film coated round brown tablets with SK on one side and 100 on the other side
150 mg	71699-150-30	Film coated round light orange tablets with SK on one side and 150 on the other side
200 mg	71699-200-30	Film coated modified oval light orange tablets with SK on one side and 200 on the other side

Titration Blister Packs; 14-Day

Daily Dose	NDC Number	Supplied As [strength(quantity)]	Tablet Description (Color, Shape, Markings)
12.5 mg per day for 14 days	71699-204-14	12.5 mg (14-count)	Uncoated round white to off-white tablets with SK on one side and 12 on the other side

Titration Blister Packs; 28-Day

Daily Dose	NDC Number	Supplied As [strength(quantity)]	Tablet Description (Color, Shape, Markings)
12.5 mg per day for 14 days, then 25 mg per day for 14 days	71699-201-28	12.5 mg (14-count)	Uncoated round white to off-white tablets with SK on one side and 12 on the other side
		25 mg (14-count)	Film coated round brown tablets with SK on one side and 25 on the other side
50 mg per day for 14 days, then 100	71699-202-28	50 mg (14-count)	Film coated round yellow tablets with SK on one side and 50 on the other side

mg per day for 14 days		100 mg (14-count)	Film coated round brown tablets with SK on one side and 100 on the other side
150 mg per day for 14 days, then 200 mg per day for 14 days	71699-203-28	150 mg (14-count)	Film coated round light orange tablets with SK on one side and 150 on the other side
		200 mg (14-count)	Film coated modified oval light orange tablets with SK on one side and 200 on the other side

Maintenance Blister Packs; 28-Day

Daily Dose	NDC Number	Supplied As [strength(quantity)]	Tablet Description (Color, Shape, Markings)
250 mg per day	71699-104-56	100 mg (28-count)	Film coated round brown tablets with SK on one side and 100 on the other side
		150 mg (28-count)	Film coated round light orange tablets with SK on one side and 150 on the other side
350 mg per day	71699-103-56	150 mg (28-count)	Film coated round light orange tablets with SK on one side and 150 on the other side
		200 mg (28-count)	Film coated modified oval light orange tablets with SK on one side and 200 on the other side

16.2 Storage and Handling

Store XCOPRI tablets at 20°C to 25°C (68°F to 77°F) with excursions permitted to 15°C to 30°C (59°F to 86°F) (*See USP Controlled Room Temperature*).

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

DRESS/Multi-organ Hypersensitivity

Instruct patients and caregivers that a fever or rash associated with signs of other organ system involvement (e.g., lymphadenopathy, hepatic dysfunction) may be drug-related and should be reported to their healthcare provider immediately. XCOPRI should be discontinued immediately if a serious hypersensitivity reaction is suspected [*see Warnings and Precautions (5.1)*].

QT Shortening

Instruct patients to inform their healthcare provider of all of the medications, over-the-counter medications, and herbal supplements that they are taking. Instruct patients to notify their healthcare provider if they have any symptoms of shortening of the QT interval, including prolonged heart palpitations or a loss of consciousness [*see Warnings and Precautions (5.2)*].

Suicidal Behavior and Ideation

Counsel patients, their caregivers, and/or families that antiepileptic drugs, including XCOPRI, may increase the risk of suicidal thoughts and behavior, and advise patients to be alert for the emergence

or worsening of symptoms of depression; unusual changes in mood or behavior; or suicidal thoughts, behavior, or thoughts about self-harm. Advise patients, their caregivers, and/or families to report behaviors of concern immediately to a healthcare provider [*see Warnings and Precautions (5.3)*].

Neurological Adverse Reactions

Counsel patients that XCOPRI causes somnolence, fatigue, dizziness, and gait disturbance. These adverse reactions, if observed, are more likely to occur early in treatment but can occur at any time. Advise patients not to drive or operate machinery until they have gained sufficient experience on XCOPRI to gauge whether it adversely affects their ability to drive or operate machinery and that other CNS depressants or alcohol may have additive effects [*see Warnings and Precautions (5.4)*].

Withdrawal of XCOPRI

Advise patients not to discontinue use of XCOPRI without consulting with their healthcare provider. XCOPRI should normally be gradually withdrawn to reduce the potential for increased seizure frequency and status epilepticus [*see Warnings and Precautions (5.5)*].

Contraceptives

Counsel females of reproductive potential that XCOPRI may decrease the efficacy of oral contraceptives and advise them to use additional or alternative non-hormonal birth control [*see Drug Interactions (7.1)*].

Pregnancy

Advise patients to notify their healthcare provider if they become pregnant or intend to become pregnant during XCOPRI therapy. Encourage patients to enroll in the North American Antiepileptic Drug Pregnancy Registry if they become pregnant. This registry is collecting information about the safety of antiepileptic drugs during pregnancy [*see Use in Specific Populations (8.1)*].

Dosing Instructions

Counsel patients that XCOPRI may be taken any time with or without food. Instruct patients that XCOPRI tablets can be taken whole or crushed. The crushed tablet can be mixed with water and either administered by mouth as an oral suspension or administered via a nasogastric (NG) tube. Counsel patients administering XCOPRI as an oral suspension or via NG tube on appropriate preparation and administration instructions [*see Dosage and Administration (2.3)*].

Abuse and Dependence

Advise patients that XCOPRI is a federally controlled substance (CV) because it can be abused or lead to dependence [*see Drug Abuse and Dependence (9)*]. Advise patients to keep their medication in a safe place to prevent misuse and abuse.

Manufactured for:
SK Life Science, Inc.
Paramus, NJ 07652

Phone: 1-866-657-5574
© 2020 SK Life Science, Inc.

XCOPRI® is a registered trademark of SK Life Science, Inc. or its affiliates and protected by U.S. Patent 7,598,279